



STATE OF CALIFORNIA  
MARINE RESEARCH COMMITTEE




**CALIFORNIA COOPERATIVE OCEANIC  
FISHERIES INVESTIGATIONS**


**ATLAS No. 14**




Thaliacian




Amphipod




Crustacean larva




Chaetognath



Copepod



Pteropod



Euphausiid

CALIFORNIA  
COOPERATIVE  
OCEANIC  
FISHERIES  
INVESTIGATIONS

*Atlas No. 14*

STATE OF CALIFORNIA  
MARINE RESEARCH COMMITTEE

*Cooperating Agencies:*  
CALIFORNIA ACADEMY OF SCIENCES  
CALIFORNIA DEPARTMENT OF FISH AND GAME  
STANFORD UNIVERSITY, HOPKINS MARINE STATION  
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION, NATIONAL MARINE FISHERIES SERVICE  
UNIVERSITY OF CALIFORNIA, SCRIPPS INSTITUTION OF OCEANOGRAPHY

June, 1971

## THE CALCOFI ATLAS SERIES

This is the fourteenth in a series of atlases containing data on the hydrography and plankton from the region of the California Current. The field work was carried out by the California Cooperative Oceanic Fisheries Investigations,<sup>1</sup> a program sponsored by the State of California under the direction of the State's Marine Research Committee. The cooperating agencies in the program are:

California Academy of Sciences  
California Department of Fish and Game  
Stanford University, Hopkins Marine Station  
National Oceanic and Atmospheric Administration, National Marine Fisheries Service<sup>2</sup>  
University of California, Scripps Institution of Oceanography

CalCOFI atlases<sup>3</sup> are issued as individual units as they become available. They provide processed physical, chemical and biological measurements of the California Current region. Each number may contain one or more contributions. A general description of the CalCOFI program with its objectives appears in the preface of Atlas No. 2.

This atlas was prepared by the Data Collection and Processing Group of the Marine Life Research Program, Scripps Institution of Oceanography.

### CalCOFI Atlas Editorial Staff:

Abraham Fleminger and Hans T. Klein, Editors  
John G. Wyllie, Cartographer

CalCOFI atlases in this series, through June 1971, are:

- No. 1. Anonymous, 1963. CalCOFI atlas of 10-meter temperatures and salinities 1949 through 1959.
- No. 2. Fleminger, A., 1964. Distributional atlas of calanoid copepods in the California Current region, Part I.
- No. 3. Alvariño, A., 1965. Distributional atlas of Chaetognatha in the California Current region.
- No. 4. Wyllie, J. G., 1966. Geostrophic flow of the California Current at the surface and at 200 meters.
- No. 5. Brinton, E., 1957. Distributional atlas of Euphausiacea (Crustacea) in the California Current region, Part I.
- No. 6. McGowan, J. A., 1967. Distributional atlas of pelagic molluscs in the California Current region.
- No. 7. Fleminger, A., 1967. Distributional atlas of calanoid copepods in the California Current region, Part II.
- No. 8. Berner, L., 1967. Distributional atlas of Thaliacea in the California Current region.
- No. 9. Kramer, D., and E. H. Ahlstrom, 1968. Distributional atlas of fish larvae in the California Current region: Northern Anchovy, *Engraulis mordax* Girard, 1951 through 1965.
- No. 10. Isaacs, J. D., A. Fleminger and J. K. Miller, 1969. Distributional atlas of zooplankton biomass in the California Current region: Spring and Fall 1955-1959.
- No. 11. Ahlstrom, E. H., 1969. Distributional atlas of fish larvae in the California Current region: jack mackerel, *Trachurus symmetricus*, and Pacific hake, *Merluccius productus*, 1951 through 1966.
- No. 12. Kramer, D., 1970. Distributional atlas of fish eggs and larvae in the California Current region: Pacific sardine, *Sardinops caerulea* (Girard), 1951 through 1966.
- No. 13. Smith, P. E., 1971. Distributional atlas of zooplankton volume in the California Current region, 1951 through 1966.
- No. 14. Isaacs, J. D., A. Fleminger and J. K. Miller, 1971. Distributional atlas of zooplankton biomass in the California Current region: Winter 1955-1959.

<sup>1</sup>Usually abbreviated CalCOFI, sometimes CALCOFI or CCOFI.

<sup>2</sup>Formerly called U. S. Fish and Wildlife Service, Bureau of Commercial Fisheries.

<sup>3</sup>For citation this issue in the series should be referred to as CalCOFI Atlas No. 14.

DISTRIBUTIONAL ATLAS OF ZOOPLANKTON BIOMASS IN THE  
CALIFORNIA CURRENT REGION: WINTER 1955-1959

J. D. Isaacs, A. Fleminger and J. K. Miller

CALCOFI ATLAS NO. 14

Data Collection and Processing Group  
Marine Life Research Program  
Scripps Institution of Oceanography  
La Jolla, California

June, 1971

# DISTRIBUTIONAL ATLAS OF ZOOPLANKTON BIOMASS IN THE CALIFORNIA CURRENT REGION: WINTER 1955-1959

J. D. Isaacs, A. Fleminger and J. K. Miller<sup>1</sup>

## Text

Introduction . . . . .	v
Materials and Methods . . . . .	vi
Sample Analyses: Procedures . . . . .	viii
Presentation of Data . . . . .	ix
Zooplankton Abundance and Diurnal Vertical Migration . . . . .	xiii
List of Charts . . . . .	xvii
References . . . . .	xviii
Appendix I . . . . .	xxi
Appendix II . . . . .	xxii

## Charts

CalCOFI Basic Station Plan . . . . .	1
All Taxa Combined . . . . .	2
Amphipoda . . . . .	8
Chætognatha . . . . .	18
Cladocera . . . . .	23
Copepoda . . . . .	28
Crustacean larvæ . . . . .	36
Ctenophora . . . . .	44
Decapoda . . . . .	49
Euphausiacea . . . . .	57
Heteropoda . . . . .	67
Larvacea . . . . .	73
Medusæ . . . . .	78
Mysidacea . . . . .	83
Ostracoda . . . . .	88
Pteropoda . . . . .	97
Radiolaria . . . . .	104
Siphonophora . . . . .	110
Thaliacea . . . . .	117

## INTRODUCTION

Beginning in 1949, the California Cooperative Oceanic Fisheries Investigations (CalCOFI) program has systematically monitored a variety of oceanographic features across the California Current region. Standardized measurements and collections of physical, chemical and biological components are taken from a network of stations. The stations are usually spaced at 40-mile intervals between 25°N and

38°N and extend seaward to several hundred miles off the Pacific coast of North America (Chart 1). The California Current region, distinguished by upwelling and relatively high productivity, comprises the eastern boundary current of the north Pacific Ocean.

The CalCOFI Zooplankton Collection has already served a large number of specialized studies concerned with distribution and abundance of organisms among the epipelagic fauna of the region. For example, data in CalCOFI Atlases 2, 3 and 5 through 9 confirm the eastern features of the biogeographical patterns proposed for the mixed layers of the north Pacific Ocean (Beklemishev, 1967; McGowan, 1971). The CalCOFI data illustrate the relatively stable relationships among these features as well as their seasonal latitudinal displacement. In addition to seasonal variation, striking distributional changes have also occurred between periods of several years duration (Ahlstrom, 1960; Balech, 1960; Berner, 1960; Berner and Reid, 1961; Brinton, 1960; Radovich, 1960). Biological fluctuations are better understood when viewed in the context of a broad environmental perspective. The CalCOFI program is providing concurrent measurements of time and space related hydrographic, meteorological and biological events. The ecological insights to emerge will help formulate guidelines for conserving while utilizing pelagic natural resources.

The CalCOFI standard and vertically stratified zooplankton samples, are also a unique source for conducting qualitative and quantitative studies into the range of natural and man induced changes on open ocean populations. The CalCOFI samples represent all sectors of the California Current region, north to south and nearshore to offshore, in all seasons and for all years from 1949 to the present. The

<sup>1</sup>Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California.

35,000 samples, preserved in formaldehyde and stored under curatorial supervision as part of the Zooplankton Collections of Scripps Institution of Oceanography, have a useful research life expectancy of probably more than 50 years. That is, the accruing collection comprises a relatively permanent, quantitatively comparable historical record of environmental conditions in the eastern North Pacific epipelagic waters as expressed in terms of its zooplankton, an archive that coincides with the years of maximum sustained increase in variety as well as the level of man's intervention into the oceans. Thus, these samples may serve the present and the future in analytical research activities involving fluctuations of local chemical substances, of oceanic conditions and of the stocks of hundreds of pelagic species or groups representing a highly diverse assemblage of trophic, morphologic and ethologic types.

Displacement volumes of the CalCOFI plankton collections are routinely measured and have been presented in the CalCOFI Atlas series (Smith, 1971). Displacement volumes afford a rapid estimate of the standing crop of plankton at minimal cost. However, without qualification as to contents, their contribution to an understanding of the region's ecosystem is limited.

Quantitative estimates of phylogenetically different groups may provide an estimate of nutritive quality of the standing crop as well as an index of trophodynamic complexity in the region. Biomass estimates made species by species, however, are a practical impossibility for large collections. A simpler procedure would be to group together species of higher taxonomic categories since it is probable that the components of such categories occupy similar or overlapping trophic levels. Species within a category would tend to be like each other in morphology, nutrient storage and size, and therefore similar in terms of nutrient quality relative to other categories. For example, chaetognaths are primarily metazoan carnivores, salps are fine-screen filterfeeders and epiplank-

tonic copepods range from coarser screen filter-feeders to obligate secondary consumers. These three taxonomically high-ranking groups not only occupy somewhat different positions in the trophic spectrum but are quite different from one another with respect to body proportions, size range of adults, proportion of water in body weight and presumably mechanisms of defense against predation. In general, therefore, taxonomic categories may also be viewed ecologically and nutritionally as "functional groups."

The number of functional groups to be employed in analyzing the distribution of biomass in a set of samples must of necessity be determined by the optimum desired resolution versus the limitations of the laboratory. In the present study, zooplankton biomass is being measured in terms of 17 functional groups (Table 1). The wet weight of each functional group corrected for interstitial water is estimated in every sample. All of the standard zooplankton collections from 20 CalCOFI cruises, namely, mid-spring, mid-summer, mid-fall and mid-winter cruises from the years 1955 through 1959, are analyzed in this manner. Contoured biomass distributions of the 17 functional groups found in the mid-spring and mid-fall sets of samples were presented in CalCOFI Atlas No. 10 (Isaacs, Fleminger and Miller, 1969). The present atlas is concerned with mid-winter cruises.

## MATERIALS AND METHODS

Selection of the years 1955 through 1959 for analysis of biomass distribution was dictated by interest in the occurrence and nature of patterns of seasonal and annual variability among the functional groups of zooplankton. During this time, yearly mean temperatures above the thermocline shifted upward from the relatively cold years of 1955 and 1956 to the relatively warm years of 1958 and 1959. These changes in the California Current region are well documented (Anon., 1963) and broader ramifications as experienced in other areas within the north Pacific were discussed in a

CalCOFI-sponsored symposium entitled "The Changing Pacific Ocean in 1957 and 1958," (Sette and Isaacs, Eds., 1960).

In the present atlas, winter is represented by the January cruises of 1955 through 1959 (Table 2). Similar analyses of the summer season, now in preparation, are scheduled to appear in a future CalCOFI atlas. The net and the procedures employed in obtaining CalCOFI plankton samples have already been described at length (Smith, 1971). Suffice it to say that the net is one meter in diameter at the mouth and about five meters long, the conical straining section having a mesh size of approximately 0.5 mm. The net, towed obliquely from a ship underway at about 2 knots, makes a traverse from the surface to 140 meters and return, requiring about 14.5 minutes and usually strain-

ing 400 to 600 cubic meters of sea water.

Analysis of samples for this atlas has been carried out by the Biomass Laboratory, a team established for this purpose within the Marine Life Research Group at Scripps Institution of Oceanography. This laboratory is regularly staffed by a small number of technically trained personnel and also employs temporary and part-time employees recruited from among the student body of the University of California at San Diego. Newcomers undergo a training phase followed by a prolonged apprenticeship. During this trial period their work is closely scrutinized and must satisfactorily meet a number of quality control tests before their analyses of samples are included among the laboratory's data output. Plankton specialists at Scripps Institution participated with the lab-

**Table 1.** Number of epiplanktonic species in CalCOFI region.

Functional Group <sup>1</sup>	Approximate number of species found in regions between the surface and 140 meters depth; source of estimate.	
Amphipoda	≥ 41	(Bowman, 1953)
Chaetognatha	24	(Alvariño, 1965)
Cladocera	5	(Baker, 1938; Strickland, et al., 1968)
Copepoda	≥ 235	(Esterly, 1905; Olson, 1949; Fleminger, 1967)
Crustacean larvae	No estimate available	
Ctenophora	≥ 4	(Torrey, 1904)
Decapoda	No estimate available	
Euphausiacea	28	(Brinton, 1967a)
Heteropoda	13	(McGowan, 1967)
Larvacea	26	(Tokioka, 1960)
Medusae	24	(Alvariño, pers. comm.)
Mysidacea	≥ 14	(Banner, 1948; Tattersall, 1951; W. D. Clarke, pers. comm., 1969)
Ostracoda	≥ 16	(Juday, 1906; C. Miller, pers. comm., 1969)
Pteropoda	29	(McGowan, 1967)
Radiolaria, Tripylea <sup>2</sup>	≥ 20	(Kling, 1966; P. B. Helms and E. D. Milow, pers. comm., 1969) <sup>2</sup>
Thaliacea	≥ 23	(Berner, 1967)
Siphonophora	44	(Alvariño, pers. comm.)
All Taxa Combined	≥ 546	

<sup>1</sup>Fish eggs, fish larvae and squid are regularly removed from CalCOFI plankton samples by the La Jolla Laboratory, Bureau of Commercial Fisheries, U.S. Fish and Wildlife Service and are being reported elsewhere. Within the functional group "Decapoda" we have included for convenience galatheid crabs, holoplanktonic Natantia, planktonic juveniles of other Natantia and planktonic juveniles of stomatopods (Hoplcarida). Our functional group "Crustacean larvae" is comprised of the pelagic larvae of Brachyura and Anomura without the galatheids. Other identifiable larvae, crustacean or otherwise, are included within their respective functional groups. In general the forms included herein are those large enough to be retained regularly by 0.5 mm mesh.

<sup>2</sup>Radiolarian counts usually do not include the acantharians, often poorly preserved, and the much smaller-sized polycystins; adding the latter two groups brings the number of radiolarian species in the California Current region to about 200 (R. Casey, pers. comm., 1969).

oratory supervisor, Mr. Julian K. Miller, in training the initial cadre of the Biomass Laboratory and subsequent recruits have been trained and tested by the Laboratory's senior personnel. The usual quality control test consists of re-submitting selected samples already analyzed but disguised by substituting a false label for the original one. Re-analysis is performed by either the same worker or some other mem-

ber of the laboratory. Selection of the operator, the sample and the frequency of the test is random unless deviations from the standards are excessive. Variation of less than  $\pm 5\%$  is regarded as an acceptable performance. The reliability of data produced in this laboratory has been tested independently by Professor E. W. Fager of Scripps Institution, who has kindly provided a summary of results in Appendix I.<sup>3</sup>

**Table 2.** CalCOFI zooplankton samples selected for analysis of winter conditions. The first two digits of the cruise number represent the year, the last two, the month of a cruise.

Cruise	Number of samples
Winter	
5501	111
5601	110
5701	57
5801	246
5901	238
	762

### SAMPLE ANALYSES: PROCEDURES

The operator analyzing the samples seeks to obtain a representative wet weight, less interstitial water, standardized to grams per 1000m<sup>3</sup> of sea water, for each functional group present in the sample. These weights comprise an estimate of standing crop occurring between the surface and about 140 m depth. Weight has been determined indirectly by means of a standardized length, as described below.

In practice, analysis begins with displacement volumes following the method of Ahlstrom and Thrailkill (1963) taken before and after removal of "large jellies" (thaliaceans, cnidarians, heteropods). Remaining jellies are sorted into their respective functional groups and weight is estimated directly with the aid of an analytical balance. The biomass of the other functional groups is determined from an aliquot ranging in size from 1/2 to 1/32, depending upon the volume of the sample after removal of the large jellies. As a rule the size of the aliquot examined is at least 1 cc in displacement volume. Its adequacy is judged by comparison with the displacement volume of the total sample as described below. The aliquot is obtained with the aid of a Folsom split-

ter (McEwen, Johnson and Folsom, 1954).

The aliquot is poured into a square transparent dish of plexiglass 20 by 20 cm whose chamber is partitioned into seven equal sub-chambers by six equidistant parallel ridges. The contents are examined with the aid of a stereomicroscope at 7X magnification. One of the oculars is fitted with a ruled ocular disc bearing 100 equally spaced divisions (1 division = 0.167 mm at 7X magnification) that is used to estimate the length of each specimen. Standard length measurements for conversion to weight (conversion tables are presented in Appendix II) are made to the nearest division. The specimens are tallied in size classes which vary somewhat depending on the overall size range of the functional group. The points of reference for taking a standard length in each functional group are shown in Figure 1. Where practical, bent specimens are straightened and, based on experience, allowances are made for slight to moderate distortions. Grossly distorted specimens are removed and weighed on a balance. To eliminate the effect of interstitial

<sup>3</sup>We thank Prof. E. W. Fager for devising and analyzing this test and Mr. G. T. Hemingway for requesting the test and for supervising the gathering of the data.

water the standard weight is corrected by reducing it to two-thirds of the gross weight (Ahlstrom and Threlkill, 1963). This correction for interstitial water has been incorporated in the length-weight conversion tables of Appendix II. Weights are also taken directly of functional groups dominated by a single growth stage or a species that is morphometrically atypical with respect to its functional group. The conversion tables are based upon typical samples in which each functional group consists of a heterogeneous mixture of species and growth stages.

The estimated weight of all functional groups within a sample are summed, and to be acceptable, must be within 10% of the normalized weight based upon displacement volume of the total sample. Use of the displacement volume to estimate weight is predicated upon the relationship of volume and weight in stabilized preserved samples, as determined by Ahlstrom and Threlkill (1963).

## PRESENTATION OF DATA

In general, practices established in earlier atlases of this series have been followed. The results of each cruise is shown separately by functional group and plotted on standard charts. We have departed from previous contouring practices by using contour intervals based upon a factor of 4. The change from our usual factor of 10 was dictated by (1) the range of the biomass values extending below as well as above unity, (2) the desirability of having the value, one gram (i.e., one part organism in  $10^9$  parts sea water) coincide with a contour line, (3) the need for a system of interpolable logarithmic contours and (4) the range (0.5 to 2 times) of the 95% confidence limits of any single estimate with respect to variability from laboratory procedures employed in sample analysis (see Appendix I).

The biomass estimates are expressed in terms of grams wet weight per  $1000\text{m}^3$  of sea water strained (corrected for interstitial water) per functional group. Contour intervals are factors of 4 and contour lines coincide with a se-

quence of values as follows: 0.016, 0.062, 0.25, 1, 4, 16, 64. The contour intervals increase exponentially and the contours are defined by the equation:

$$C_n = KB^{an}$$


where

$C_n$  is the value of a contour for a value of the exponent  $n$ ,  
 $B$  is the base,  
 $K$  is a constant,  
 $a$  is constant, and  
 $n$  is a variable with integer values.


As employed here,  $K = 1$ ,  $B = 4$ ,  $a = 1$ , and  $n = -3, -2, -1, 0, 1, 2, 3$ , etc. The system permits indefinite extension above and below unity. Interpolation or extrapolation into a finer or coarser series of logarithmic intervals can be readily accommodated.

No more than seven contour intervals are shown per chart and seven grades of shading are applied in the same sequence of increasing abundance. The sequence of seven values within the exponential series used on a specific chart is the one that best fits the range of abundance estimates shown by a functional group in a specific cruise. The highest and the lowest estimates of biomass (greater than zero) are shown in Arabic numerals to the right alongside the index of contour interval shadings in the chart legend. The charts have been arranged by functional groups which follow an alphabetical sequence, the sum total of the functional groups appearing first under the heading "All Taxa Combined." Within each category the charts are arranged by season, each seasonal set of five cruises following in chronological order.


In addition to the contoured estimates of biomass overlying the station plan, each chart presents two graphic summaries pertaining to the abundance of the functional group (Fig. 2). Latitudinal distribution of biomass is estimated in terms of the mean weight per station in each set of three successive lines of stations. This is shown as a histogram to the left of the distribution. The proportion of the functional group to the category All Taxa Combined, i.e.,




*Thaliacea*




*Copepoda*




*Amphipoda*




*Ctenophora*




*Euphausiacea*



*Mysidacea*




*Radiolaria*





*Heteropoda*

**Figure 1.** Standard lengths used in this study. The measurements delimited by arrows are taken with the aid of an ocular micrometer at 7X magnification.



**Figure 1, cont.** Tables showing the relationship of standard length to weight among zooplankton from the California Current region appear in Appendix II.



**Figure 2.** An example of the charts presented in this Atlas. Special quantitative summaries include: 1. ratio of quantity caught in night samples relative to that in twilight and day samples; 2. north to south comparison of abundance; 3. total amount of the functional group taken relative to that of the entire catch (All Taxa Combined).

the total zooplankton biomass per cruise less fish eggs, fish larvae and squid is shown in actual values and in percentage in a bar diagram located in the lower left margin of the chart.

## ZOOPLANKTON ABUNDANCE AND DIURNAL VERTICAL MIGRATION

CalCOFI monitoring cruises usually survey large areas exceeding 200,000 square miles and occupy 200 or more stations during a 30 to 40-day period. The severe time limitations imposed upon the sampling program do not permit the sampling of zooplankton with regard to time of day, intensity and quality of light striking the sea surface or water transparency. It is well known that samples of zooplankton from the mixed layer vary appreciably in both quality and quantity when collected under similar conditions but at different times within a 24-hour cycle (King and Demond, 1963; King and Hida, 1954, 1957a, b). This phenomenon is known to prevail in the California Current region (e.g., Kramer, 1963), and Brinton (1962, 1967) has shown that species of euphausiids may vary in abundance by an order of magnitude or more in replicate samples differing primarily in the hour of collection.

Seasonal influences on biotic and abiotic factors, extent of cloud cover, moon phase, water transparency, etc., may influence in part the abundance and quality of day and night samples of zooplankton. At present, however, the level of understanding of horizontal and vertical distribution in local waters is inadequate to standardize abundance and quality of samples taken at different times within a 24-hour cycle.

Some of the functional groups considered here fail to show consistent day-night differences. Moreover, among the highly diverse groups such as copepods some taxa do not ordinarily migrate below the sampling level (e.g., *Clausocalanus*, most candaciids, pontellids), others may virtually disappear from the uppermost 140 meters in daylight (e.g., *Pleuromamma*, *Undeuchaeta*) while still others may be

predominantly epipelagic in one portion of the region and mesopelagic when present elsewhere (e.g., *Calanus*, *Eucalanus*, *Rhincalanus*).

We have provided three visual aids to assist in recognizing possible artifacts in a distribution introduced by time of sampling. Each station symbol shows whether the sample was obtained during the hours of daylight, twilight (sunrise or sunset) or night. Twilight is used here to denote a period of time extending one-and-one-half hours before and one-and-one-half hours after local sunrise or local sunset. The terms day and night are restricted to the hours between the two twilight periods.

Ratios of the median values of night (N), twilight (T) and Day (D) catches are shown in the chart legend as N/D and N/T. When these ratios are close to unity it is assumed that the contour intervals absorb much if not all of the variability due to time of sampling. The relationship between these ratios and the distribution of day, twilight and night values for a few typical examples is shown in Figure 3.


The less abundant functional groups were frequently absent from one-half or more of the total number of observations per cruise. In these instances, N/D and N/T ratios were estimated graphically. Abundance was ranked by cruise and rank was plotted against abundance. To normalize differences in the number of observations between time periods under comparison, the linear distance between units along the scale of ranks for the set of fewer observations was adjusted to match the distance required for the ranks of the set with the higher number of observations. If the two resulting curves are roughly parallel the N/D and N/T ratios are determined by the distance between the two lines along the ordinate scale (Fig. 3). If the curves were strongly oblique or crossing, the differences were assumed to be due to factors other than time of day and N/D or N/T ratios were not estimated.

Contoured distributions may be noticeably influenced by the time of sampling when the ratio of the median exceeds 2. Accordingly, to compensate for artifacts when N/D or N/T

equalled or exceeded 2, an additional chart was prepared in which day and twilight catches were multiplied by N/D and N/T, respectively, so as to more closely approximate an equivalent night-time distribution. Each of the adjusted charts is preceded by a contoured presentation of the unfactored data.

Variability due to the hour of collection appears to have a consistent and noticeable effect on the charted distributions in only certain of the functional groups, namely euphausiids, am-

phipods and perhaps copepods. Adjustment of the estimated abundances in these cases, however, has resulted in only moderately improved coherence within the distributions. Due to the general agreement among the large number of samples in our data and evidence based on repeated sampling of euphausiids (Brinton, 1962; Figs. 16-23), we believe the charts reflect real geographical variations in relative abundance, irrespective of accumulated errors from escapement, avoidance, patchiness and



**Figure 3.** Scatter diagram of three representative examples showing the distribution of biomass ranked in order of increasingly higher values. Each set of data consists of all measurements of a functional group derived from a single cruise. Day, twilight and night samples are differentiated and the scale of ranked position (abscissa) is adjusted so that the number of samples per time period extends over the same length irrespective of the number of observations. Further comments in the text.

vertical migration.

Questions arose over the use of medians derived from unpaired sets of observations to determine the ratios, N/D and N/T. Paired observations may be simulated by pairing samples from adjacent stations along the cruise track which were occupied successively in different time phases (day, twilight or night). Pairing

increases the likelihood that differences in the catch reflect variability due primarily to time of collection; i.e., the sets of observations were made on ecologically and biogeographically similar assemblages of organisms. Comparisons between the ratios of such simulated pairs with unpaired data were carried out for various representative functional groups (Table 3).

Table 3. N/D ratios for paired and unpaired observations.


	Unpaired observations		Paired observations
	mean N mean D	median N median D	Median $\frac{N}{D}$
<b>Cruise 5510</b>			
All Taxa Combined	1.5	1.9	1.9
Euphausiacea	4.0	6.5	6.8
Copepoda	1.5	2.0	2.0
Pteropoda	1.9	3.8	2.4
Ostracoda	3.5	6.7	8.0
<b>Cruise 5910</b>			
All Taxa Combined	1.0	1.4	1.5

The ratios of paired observations were found to scatter broadly. No more than about one-third were within  $\pm 0.5$  of either the ratio of the medians or the general trend observable in graphed data similar to Figure 3. On the other hand, the median ratio of the paired sets (Table 3) is remarkably similar to the ratio of the medians of unpaired observations as well as to the trend of the differences between the day and night curves in Figure 3. We conclude that simulating paired observations does not noticeably improve upon the use of the ratio of the medians, N/D and N/T, in compensating for day, twilight and night effects on standing-crop estimates from standard CalCOFI tows.

Our method for adjusting catch size relative to time of sampling was chosen over the sine curve method employed on mid-Pacific plankton by King and Hida (1954, 1957a, b) for several reasons. Chief among them are: (1) our practice of estimating biomass separately for each functional group; (2) our region of study which spans a biogeographical gradient from the boreal to the tropical and an ecological gradient from coastal-neritic water to cen-

tral waters overlying the ocean basin; (3) the full range of variability shown by our samples with respect to their individual extent of faunal homogeneity. We are reluctant to assume that different combinations of faunally heterogeneous species comprising a functional group will tend to behave diurnally in a similar manner. In fact, N/D and N/T may show considerable variability within functional groups (Fig. 3) and the ratio of the medians differs appreciably between groups. It might also be noted that compared to the sine curve method, calculation and employment of the ratio of the medians was carried out more easily.

Using the data for All Taxa Combined from Cruise 5904 and for euphausiids, from Cruise 5704, both selected solely for the sake of convenience, direct comparisons of contour patterns were made between values adjusted for differences in the time of catch by means of the King and Hida sine curve factor and the ratio of the medians factor. With N/D being 2.04 for All Taxa Combined and 24.5 for the euphausiid data, the two examples approximate the breadth of the range in the ratio of the medians encountered during the study. We



**Figure 4.** Comparison of two methods to compensate for the effect of diurnal vertical migration on plankton abundance. Categories "All Taxa Combined" and "Euphausiacea" were selected as examples to show moderate and pronounced differences in abundance, respectively, among samples taken at different times of the 24-hour cycle. In B, C and D the abundance of samples taken in hours other than the desired sampling time has been adjusted by the appropriate factor. The distribution of the biomass estimates is contoured in the usual manner.

(A) unadjusted values; (B) day and twilight values adjusted by the ratio of the medians method to approximate median night values; (C) values adjusted by the sine curve method (King and Hida, 1954) to approximate abundance at 0000 hours; (D) as in C, above, but adjusted to approximate abundance at 0600 and 1800 hours when sine is zero as originally employed by King and Hida (1954). Additional explanation and comments in text.

found no noticeable differences in either the general pattern or in coherence of the contours between the sets of data adjusted by the two different procedures (Fig. 4).

It should be noted that in contrast to the adjustments by King and Hida, we have normalized data to represent median night values,

typically the period of time yielding the highest standing crop for most of the functional groups. Computer programs for adjusting catch size by the sine curve method or by the ratio of the medians method are on file at Scripps Institution of Oceanography. A copy may be obtained by writing Prof. E. W. Fager.

**Table 4.** List of charts.

Cruise Number	Winter				
	5501	5601	5701	5801	5901
All Taxa Combined	2	3	4	5	6
All Taxa Combined, adjusted					7
Amphipoda	8	10	12	14	16
Amphipoda, adjusted	9	11	13	15	17
Chaetognatha	18	19	20	21	22
Cladocera	23	24	25	26	27
Copepoda	28	30	31	32	34
Copepoda, adjusted	29			33	35
Crustacea larvae	36	38	39	41	43
Crustacea larvae, adjusted	37		40	42	
Ctenophora	44	45	46	47	48
Decapoda	49	51	53	54	55
Decapoda, adjusted	50	52			56
Euphausiacea	57	59	61	63	65
Euphausiacea, adjusted	58	60	62	64	66
Heteropoda	67	68	69	70	71
Heteropoda, adjusted					72
Larvacea	73	74	75	76	77
Medusae	78	79	80	81	82
Mysidacea	83	84	85	86	87
Ostracoda	88	89	91	93	95
Ostracoda, adjusted			90	92	96
Pteropoda	97	98	100	101	102
Pteropoda, adjusted			99		103
Radiolaria	104	106	107	108	109
Radiolaria, adjusted	105				
Siphonophora	110	111	113	114	115
Siphonophora, adjusted			112		116
Thaliacea	117	118	119	120	121
Thaliacea, adjusted					122

## REFERENCES CITED

- Ahlstrom, E. H., 1948. A record of pilchard eggs and larvæ collected during surveys made in 1939 to 1941. U. S. Fish Wildl. Serv. Spec. Sci. Rep., Fish. No. 54: 76 pp.
- \_\_\_\_\_, 1952. Pilchard eggs and larvæ and other fish larvæ, Pacific Coast, 1950. U. S. Fish Wildl. Serv. Spec. Sci. Rep., Fish No. 80: 58 pp.
- \_\_\_\_\_, 1954. Distribution and abundance of egg and larval populations of the Pacific sardine. U. S. Fish Wildl. Serv., Fish. Bull., 93(56): 83-140.
- \_\_\_\_\_, 1960. Fish spawning in 1957 and 1958. CalCOFI Repts., 7: 173-179.
- \_\_\_\_\_, and J. R. Thraillkill, 1963. Plankton volume loss with time of preservation. CalCOFI Repts., 9: 57-73.
- Alvariño, A., 1965. Distributional atlas of Chaetognatha in the California Current region. CalCOFI Atlas No. 3: vii-xiii, 1-291.
- Anonymous, 1963. CalCOFI atlas of 10-meter temperatures and salinities, 1949 through 1959. CalCOFI Atlas No. 1; iii-iv, 1-297.
- Baker, H. M., 1938. Studies on the Cladocera of Monterey Bay. Proc. Calif. Acad. Sci., 4th Ser., XXIII: 311-365.
- Balech, E., 1960. The changes in the phytoplankton population off the California coast. CalCOFI Repts., 7: 127-132.
- Banner, A. H., 1948. A taxonomic study of the Mysidacea and Euphausiacea (Crustacea) of the northeastern Pacific. Part I. Mysidacea from the family Lophogastridae through tribe Erythropini. Trans. Royal Canadian Inst., 26: 345-399.
- Beklemishev, K. V., 1967. Biogeographical division of the pelagic portion of the Pacific Ocean. In The Pacific Ocean, Vol. VII, Part I. — Plankton. Ed., V. G. Bogorov. Moscow. Trans. 435, U. S. Naval Oceanographic Office, 1969: 100-269.
- Berner, L. D., 1960. Unusual features in the distribution of pelagic tunicates in 1957 and 1958. CalCOFI Repts., 7: 133-135.
- \_\_\_\_\_, 1967. Distributional atlas of Thaliacea in the California Current region. CalCOFI Atlas No. 8: vii-xi, 1-322.
- \_\_\_\_\_, and J. L. Reid, Jr., 1961. On the response to changing temperature of the temperature-limited plankter *Doliolum denticulatum* Quoy and Gaimard, 1835. Limnol. Oceanogr., 6(2): 205-215.
- Bowman, T. E., 1953. The systematics and distribution of pelagic amphipods of the families Vibiliidae, Paraphronomidae, Hyperiidae, Dairellidae and Phrosinidae from the northeastern Pacific. Unpublished Ph.D. thesis, Univ. Calif., Los Angeles.
- Brinton, E., 1960. Changes in the distribution of euphausiid crustaceans in the region of the California Current. CalCOFI Repts., 7: 137-147.
- \_\_\_\_\_, 1962. Variable factors affecting the apparent range and estimated concentration of euphausiids in the North Pacific. Pac. Sci., 16(4): 374-408.
- \_\_\_\_\_, 1967a. Distributional atlas of Euphausiacea (Crustacea) in the California Current region, Part I. CalCOFI Atlas No. 5: vii-xi, 1-275.
- \_\_\_\_\_, 1967b. Vertical migration and avoidance-capability of euphausiids in the California Current. Limnol. Oceanogr., 12(3): 451-483.
- Esterly, C. O., 1905. The pelagic Copepoda of the San Diego region. Univ. Calif. Publ. Zool., 2(4): 113-233.
- Fleminger, A., 1964. Distributional atlas of calanoid copepods in the California Current region, Part I. CalCOFI Atlas No. 2; ix-xvi, 1-313.
- \_\_\_\_\_, 1967. Distributional atlas of calanoid copepods in the California Current region. Part II. CalCOFI Atlas No. 7: vii-xvi, 1-213.
- Isaacs, J. D., A. Fleminger and J. K. Miller, 1969. Distributional atlas of zooplankton biomass in the California Current region: Spring and Fall 1955-1959. CalCOFI Atlas No. 10: v-xxv, 1-252.
- Juday, C., 1906. Ostracoda of the San Diego region. I. Halocypridae. Univ. Calif. Publ. Zool., 3(2): 13-38.
- King, J. E., and J. Demond, 1953. Zooplankton abundance in the central Pacific. U. S. Fish Wildl. Serv., Fish. Bull., 82(54): 111-144.
- \_\_\_\_\_, and T. S. Hida, 1954. Variations in zooplankton abundance in Hawaiian waters, 1950-52. U. S. Fish Wildl. Serv. Spec. Sci. Rept., Fish. No. 118: 66 pp.

- \_\_\_\_\_, and \_\_\_\_\_, 1957a. Zoo-plankton abundance in Hawaiian waters, 1953-54. U. S. Fish Wildl. Serv. Spec. Sci. Rept., Fish. No. 221: 23 pp.
- \_\_\_\_\_, and \_\_\_\_\_, 1957b. Zoo-plankton abundance in the central Pacific. Part II. U. S. Fish Wildl. Serv., Fish. Bull., 57(118): 365-395.
- Kling, S. A., 1966. Castanellid and circoporid radiolarians: systematics and zoogeography in the eastern North Pacific. Unpublished Ph.D. thesis. Univ. Calif., San Diego.
- Kramer, D., 1963. Records and observations from plankton grid studies off Baja California, April 1952. U. S. Fish Wildl. Serv. Spec. Sci. Rept., Fish. No. 422: 42 pp.
- McEwen, G. F., M. W. Johnson and T. R. Folsom, 1954. A statistical analysis of the performance of the Folsom Plankton Sample Splitter, based on test observations. Arch. f. Meteor., Geophys. Bioklim., Ser. A: Meteor. and Geophys., Bd. 7: 502-527.
- McGowan, J. A., 1967. Distributional atlas of pelagic molluses in the California Current region. CalCOFI Atlas No. 6: vii-xiv, 1-218.
- \_\_\_\_\_, 1971. Oceanic biogeography of the Pacific. In: The Micropaleontology of Oceans, B. M. Funnell and W. R. Riedel, eds. Cambridge Univ. Press: 3-74.
- Olson, J. B., 1949. The pelagic cyclopoid copepods of the coastal waters of Oregon, California and Lower California. Unpublished Ph.D. thesis, Univ. Calif., Los Angeles.
- Radovich, J., 1960. Redistribution of fishes in the eastern North Pacific Ocean in 1957 and 1958. CalCOFI Repts., 7: 163-171.
- Sette, O. E., and J. D. Isaacs, Eds., 1960. The changing Pacific Ocean in 1957 and 1958. CalCOFI Repts., 7: 17-217.
- Smith, P. E., 1971. Distributional atlas of zooplankton volume in the California Current region, 1951-1966.
- Strickland, J. D. H., et. al., 1968. Research on the marine food chain. Progress report. January 1967-April 1968. Part IV. Data record. Coastal plankton survey 1967. Section II. The plankton. SIO Ref. 68-38. Unpublished manuscript.
- Tattersall, W. M., 1951. A review of the Mysidacea of the United States National Museum. U.S.N.M., Bull. 201: 292 pp.
- Tokioka, T., 1960. Studies on the distribution of appendicularians and some thaliaceans of the North Pacific, with some morphological notes. Publ. Seto Mar. Biol. Lab., 8(2): 351-443.
- Torrey, H. B., 1904. The ctenophores of the San Diego region. Univ. Calif. Publ. Zool., 2(2): 45-51.

## APPENDIX I<sup>1</sup>

### Reliability of Laboratory Measurements

There are a number of possible sources of error in making the counts involved in this program: incorrect splitting of samples to obtain the aliquots, operator bias, differences in efficiency in counting different classes of organisms and differences in efficiency associated with differences in relative numbers.

The first two have been examined by having replicate counts done on the same sample. In five cases the same operator made the counts but was unaware the same sample was being recounted; in 17 cases, different operators counted the same sample and the second person did not know the sample had already been counted. The median number of categories counted in each sample was 12 (range: 9-16). The signed-ranks test was applied to the paired counts on each of the samples. All five of the cases involving two counts by the same person and 15 of the 17 involving two operators were non-significant at the 20% level. This probably represents random error because it is just over the expected number at this level, namely 1.1.

Six operators were involved in the preceding test. Examination of the paired counts in the 17 cases where two persons did the counts revealed only one operator whose counts were consistently below or above those of all others who counted the same sample. This operator, D, recorded counts that were on the average somewhat below those of her partners in 7 of 8 cases ( $p = 0.10$ ). In no case was the operator sufficiently far below the partner to make the difference significant by the signed-ranks test. From these tests one can conclude that bias in sample splitting or in counting by the operators is no greater than would be expected as a result of random errors.

In order to determine whether the type of organism counted or its relative abundance had an effect on the counts, samples representing high, medium

and low abundances of 10 classes of organisms were selected. Three replicate counts were done on each sample in such a manner that the operators did not know whether a sample had been counted before or not. In addition to the counts, weights were determined in the standard fashion. An analysis of variance was run on both sets of data. The average counts for the different classes of organisms ranged from 25 to 5532; the corresponding weights ranged from 0.029 to 1.924 grams. The average counts for the three abundance classes of samples were 2623, 1026 and 47; the corresponding weights were 1.035, 0.864 and 0.047 grams. As was expected, mean squares for both of these factors and for their interaction were highly significant.

Only one of the total of six mean squares for replicate, type of organism, replicate interaction and abundance-replicate interaction was significant and this was at the 20% level. From these tests it may be concluded that neither the type of organism nor the relative abundance of the organisms have any significant effect on either the counts or the weights. The error mean square can, therefore, be used to set confidence limits for the observations. In order to make these most useful, the analyses were rerun with the data transformed to logarithms, 95% confidence limits were calculated and then transformed to anti-logarithms. The results indicate that conservative 95% confidence limits for both counts and weights are given by 0.45 and 2.20 times the observed value. These account only for the variability introduced in the laboratory handling of the samples and do not include any that may have been introduced by the sampling procedures in the field.

<sup>1</sup>Prepared by E. W. Fager, Scripps Institution of Oceanography.


**APPENDIX II<sup>1</sup>**  
**Relationship of Length to Weight, California Current Region**

Oc. Micro. Units <sup>2</sup>	No./gm	Oc. Micro. Units	No./gm	Oc. Micro. Units	No./gm
<b>Amphipods</b>					
10	1023	230	16	7	2762
20	376	240	14	10	949
30	159	250	13	13	500
40	101	260	11	15	200
50	64	270	9	25	65
60	48	280	6		
70	29	290	5	(c) Megalopa	
80	26			15	500
90	14	<b>Cladocerans</b>		20	235
100	10	3	11731	30	105
110	7	6	3921	40	68
				50	39
<b>Chaetognaths</b>		<b>Copepods</b>		60	23
20	6994	5	5814		
30	3808	10	1873	<b>Ctenophores</b>	
40	2168	15	698	Usually weighed	
50	1522	20	254		
60	762	25	123	<b>Decapods</b>	
70	614	30	70	(a) Galatheids	
80	425	35	50	30	286
90	330	40	31	40	125
100	253			50	36
110	200	<b>Crustacean larvae</b>		60	28
120	142	(a) Anomuran Zoea		70	18
130	110	3	752	80	15
140	75	5	254	90	10
150	61	10	162		
160	44	15	146	(b) Hoplocarids	
170	40	20	128	8	1368
180	34	25	111	20	775
190	31	30	100	30	253
200	27			40	217
210	23	(b) Brachyuran Zoea		50	157
220	18	4	3403	60	83

<sup>1</sup>This table was prepared by J. K. Miller and is based upon 1-gram samples of each size category of a functional group sorted from a variety of plankton collections characterized by pronounced diversity in species and in developmental stages.

<sup>2</sup>One Ocular Micrometer Unit = 0.167 mm at 7X magnification.

Oc. Micro. Units	No./gm	Oc. Micro. Units	No./gm	Oc. Micro. Units	No./gm
70	71	100	21		
80	50	110	17	15	5400
90	40	120	13	25	3420
		130	10	35	2500
(c) Natantians		140	8	45	1075
5	7143	150	7	55	625
15	1744	160	6	65	300
25	765	170	5	75	154
35	539	210	4	85	133
45	379	240	3	95	48
55	147	270	2	100	15
65	54	300	1		
75	42	330	1		
85	29	360	1	5	3465
95	20	390	1	10	1496
105	18			15	501
115	15			20	341
125	13	5	7000	25	96
135	11	10	2147	30	87
145	8	15	941		
155	7	20	525		
165	6	30	375	5	3520
175	5	40	222	10	1444
185	4	50	185	15	910
195	3	60	158	20	327
205	2	70	105	25	185
215	1.5	80	83	30	140
250	1	90	61	35	122
		100	48	40	100
<b>Euphausiids</b>		Larger forms usually weighed		45	62
5	11111			50	42
10	6243			55	33
15	4200				
<b>Larvaceans</b>					
20	1426	10-15	13200		
25	930	15-20	7173	4	5106
30	778	20-25	5200	5	3998
35	451	25-30	3848	6	2431
40	308	30-35	2564		
50	175	35-40	2197		
55	125	40-50	1923		
60	99	50-60	405		
65	89	60-70	267		
70	60				
80	45				
90	31				
		<b>Medusae</b>			
		Usually weighed			



All Taxa combined



Amphipoda



Chaetognatha



Cladocera



Copepoda



Crustacean larvae



Ctenophora



Decapoda



Euphausiacea



Heteropoda



Larvacea



Medusae



Mysidacea



Ostracoda



Pteropoda



Radiolaria





Siphonophora




Thaliacea












Biomass  
All Taxa Combined  
5701




Biomass  
All Taxa Combined  
5801




Biomass  
**All Taxa Combined**  
**5901**





Biomass  
All Taxa Combined, adjusted  
5901







Biomass  
 Amphipoda, adjusted  
 5501







Biomass  
Amphipoda, adjusted  
5601




Biomass  
 Amphipoda  
 5701




Biomass  
**Amphipoda, adjusted**  
**5701**







Biomass  
Amphipoda, adjusted


5801







Biomass  
**Amphipoda, adjusted**  
**5901**














Biomass  
Chaetognatha  
5801

























Biomass  
 Copepoda, adjusted  
 5501
























Biomass  
 Crustacean larvae, adjusted  
 5501










Biomass  
Crustacean larvae, adjusted  
5701






Biomass  
Crustacean larvae, adjusted  
580I










0.5                  %                  100  
 58                  grams              12648



Biomass  
 Ctenophora  
 5601






Biomass  
Ctenophora  
5801







Biomass  
**Decapoda**  
 5501




Biomass  
**Decapoda, adjusted**  
 5501






Biomass  
Decapoda, adjusted  
5601





Biomass  
 Decapoda  
 5701

2                  %                  100  
 grams                  2308


47










Biomass  
Decapoda, adjusted  
5901







Biomass  
**Euphausiacea, adjusted**  
 5501







Biomass  
**Euphausiacea, adjusted**  
**5601**







Biomass  
**Euphausiacea, adjusted**  
**5701**







Biomass  
Euphausiacea, adjusted  
580I








Biomass  
**Euphausiacea, adjusted**  
**5901**





















1              %              100  
72              grams              5147


Biomass  
Larvacea  
5501





















3                  %                  100  
61                  grams                  2308


Biomass  
Medusae  
5701






















0.7                  %  
 34                  grams                  5147


Biomass  
 Ostracoda  
 5501




Biomass  
Ostracoda  
5601





Biomass  
 Ostracoda, adjusted  
 5601







Biomass  
**Ostracoda, adjusted**  
 5701







Biomass  
Ostracoda, adjusted  
5801








Biomass  
Ostracoda, adjusted  
5901




















Biomass


**Pteropoda, adjusted**


5901


























Biomass


**Siphonophora, adjusted**

5601














Biomass  
**Siphonophora, adjusted**  
 5901


















These maps are designed to show essential details of the area most intensively studied by the California Cooperative Oceanic Fisheries Investigations. This is approximately the same area as is shown in color on the front cover. Geographical place names are those most commonly used in the various publications emerging from the research. The cardinal station lines extending southwestward from the coast are shown. They are 120 miles apart. Additional lines are utilized as needed and can be as closely spaced as 12 miles apart and still have individual numbers. The stations along the lines are numbered with respect to the station 60 line, the numbers increasing to the west and decreasing to the east. Most of them are 40 miles apart, and are numbered in groups of 10. This permits adding stations as close as 4 miles apart as needed. An example of the usual identification is 120, 65. This station is on line 120, 20 nautical miles southwest of station 60.

The projection of the front cover is Lambert's Azimuthal Equal Area Projection. The detail maps are a Mercator projection.

## CONTENTS

J. D. Isaacs, A. Fleminger and J. K. Miller	
Distributional atlas of zooplankton biomass in the	v
California Current region: Winter 1955 - 1959	
Charts	1-122